Eood Chemistry 41 (1991) 309-339



# Carotenoid Composition and Content of Malaysian Vegetables and Fruits by the AOAC and HPLC Methods

# E-Siong Tee

Division of Human Nutrition, Institute for Medical Research. Jalan Pahang, 50588 Kuala Lumpur, Malaysia

#### &

## Chin-Lam Lim

Food Technology Programme, School of Industrial Technology, Universiti Sains Malaysia, 11800 Pulau Pinang, Malaysia

(Received 10 August 1990; accepted 1 October 1990)

#### ABSTRACT

The *B*-carotene contents of forty vegetables and fourteen fruits were determined using the AOAC open-column (magnesia and Hyflo Super Cel mixture) chromatographic method and compared with a newly developed reverse-phase HPLC method, in which carotenoids were separated isocratically on an octadecylsilane ( $C_{18}$ ) column using a ternary mixture of acetonitrile, methanol and ethyl acetate. Results obtained showed that the AOAC method gave falsely elevated results for samples containing xcarotene, as well as those with very low  $\beta$ -carotene concentrations. On the other hand, the HPLC method successfully separated and quantitated the major carotenoids present; namely, lutein, cryptoxanthin, lycopene,  $\gamma$ - and  $\alpha$ carotenes in addition to  $\beta$ -carotene. The carotenoid composition of most of the green vegetables was rather consistent, comprising only lutein and  $\beta$ carotene. In contrast, there was no clear pattern of carotenoids present in the other vegetables and fruits, where several other carotenoids were detected m varying proportions. The vitamin A activity, expressed as  $\mu g$  of retinol equivalent (RE), was calculated on the basis of all pro-vitamin A carotenoids (cryptoxanthin,  $\gamma$ -,  $\alpha$ - and  $\beta$ -carotenes) detected. Most of the green leafy vegetables, including several local vegetables, had high RE. Several green non-leafy and other vegetables were found to have low and medium RE. None of the fruits studied may be said to have high vitamin A activity. RE calculated

309

Food Chemistry 0308-8146/91/\$03.50 (c 1991 Elsevier Science Publishers Ltd, England, Printed in Great Britain

on the basis of results from the AOAC method was found to be erroneously low for samples with significant proportions of pro-vitamin A carotenoids other than  $\beta$ -carotene, and falsely elevated for those with  $\alpha$ -carotene. Total carotenoid concentrations can be estimated by taking absorbance readings of sample extracts directly in a spectrophotometer or by the HPLC method.

The study clearly shows that the HPLC method would give a more complete picture of the carotenoid composition as well as a more accurate quantitation of the vitamin A value of the vegetables and fruits. The nutritional significance of the findings is obvious since these foods are important sources of vitamin A for the majority of the communities in the country.

#### INTRODUCTION

Vitamin A deficiency remains one of the major public health nutritional problems in many developing countries, and is an important cause of preventable blindness. The compounds pertinent in this deficiency problem, that afflicts particularly young children, are carotenoids, the major contributors of vitamin A in the diet of most communities in the region. Although an extensive literature on these compounds has been built up, carotenoids are still being actively studied all over the world, since many gaps in knowledge exist and new frontiers are being pursued (Tee, 1988).

A basic need in carotenoid research and development is knowledge of the content and composition of this group of pigments, which are widely distributed in nature and found without exception in photosynthetic tissues. In recent years, there has been particular emphasis on an understanding of the types and concentrations of various carotenoids in foods for two main reasons. First, this is of importance in relation to the pro-vitamin A activity of the carotenoids. It is thought that previously reported values of vitamin A activity in food composition tables may have been unreliable since methodologies used were not sufficiently discriminative and thus had included carotenoids that do not possess vitamin A activity (Zakaria et al., 1979; Beecher & Khachik, 1984; Underwood, 1984; Bureau & Bushway, 1986). Some of these carotenoids may occur in higher concentrations than  $\beta$ carotene, the most potent precursor of vitamin A. Second, carotenoids, including those without vitamin A activity, are now thought to play important roles beyond their classical functions in nutrition and vision. With their highly conjugated double bonds, carotenoids may act as free radical traps or antioxidants, and therefore play important roles in cancer causation and prevention (Peto et al., 1981; Olson, 1986; Temple & Basu, 1988).

The authors have embarked on a systematic study to develop improved

methodologies for the separation and quantitation of retinol and several carotenoids in foods and biological specimens, especially blood serum. A systematic review of the literature on methods for carotenoid analysis was first carried out (Tee & Lim, 1991). Initially, several studies into the physico-chemical characteristics of retinol and several carotenoids in various solvents and laboratory conditions were carried out to assist in the choice of analytical conditions. A combination of UV–vis spectroscopy and HPLC was used in these studies. An HPLC system for the analysis of these compounds was next developed, studying particularly the solvent system, peak detection and quantitation, and sample preparation. The objective was to develop a simple system, workable for routine determination of retinol and several carotenoids in a wide variety of foods of both plant and animal origin. It is hoped that the method developed could be used to re-evaluate the vitamin A value of foods tabulated in the current Malaysian Food Composition Table (Tee *et al.*, 1988).

The HPLC method developed was applied to a variety of foods. Each food sample was also simultaneously determined for vitamin A and carotenoids by the open-column chromatographic procedure of the AOAC (Williams, 1984), with the aim of determining whether the difference in  $\beta$ -carotene values thus far reported using the AOAC method were significantly different from the more specific HPLC method. If differences exist, it is hoped that the study will indicate which types of vegetables and fruits show the greatest differences. Such differences have been mentioned by various investigators, but there has not been a detailed comparison for a wide variety of foods as in the present study.

# MATERIALS AND METHODS

### Solvents and carotenoid standards

Solvents used for sample preparation and pre-treatment and for opencolumn chromatography procedures, were all analytical-grade reagents. Solvents for high-pressure liquid chromatography were of HPLC grade. All solvents for use as the mobile phase in HPLC were filtered through a 0.45  $\mu$ m regenerated cellulose membrane filter and degassed using an ultra-sonic bath.

 $\alpha$ - and  $\beta$ -Carotenes and lycopene standards were purchased from Sigma Chemical Company.  $\gamma$ -Carotene,  $\beta$ -apo-carotenal, cryptoxanthin, zeaxanthin, and lutein were gifts from F. Hoffmann La-Roche, Switzerland. The eight carotenoids used have varying structures, including the acyclic conjugated polyenelycopene: carotenoids with *psi*, *beta* and *epsilon* endgroups: oxygenated carotenoids; and an *apo*-carotenoid (Fig. 1). UV-vis



l. Lycopene (ψ, ψ-carotene)



II.  $\gamma$ -Carotene ( $\beta$ ,  $\psi$ -carotene)



III.  $\alpha$ -Carotene ( $\beta$ ,  $\epsilon$ -carotene)



IV. β-Carotene (β, β-carotene)



V. β-Cryptoxanthin; 3-hydroxy-β-carotene (β, β-caroten-3-ol)



VI. Zeaxarthin ( $\beta$ ,  $\beta$ -carotene-3,3'-diol)



VII. Lutein; xanthophyll; 3,3'-dihydroxy- $\alpha$ -carotene ( $\beta$ . $\epsilon$ -carotene-3,3'-diol)



VIII. β-apo-8'-Carotenal (8'-apo-β-caroten-8'-al)

Fig. 1. Structures of carotenoid standards (systematic names of carotenoids given in parentheses).

absorption spectra of these standards were determined and are given in Fig. 2. Stock solutions of these carotenoids were prepared in hexane (except that lutein and zeaxanthin were prepared in ethanol and  $\beta$ -apo-carotenal in petroleum ether) in concentrations of 100 µg per ml and stored in amber bottles below -20 C. Working solutions of 1 µg per ml of the standards were prepared daily. The appropriate extinction coefficients published in the literature (De Ritter, 1981) were used to calculate the exact concentration of each of the carotenoids. The preparation of all standard carotenoids was carried out with no unnecessary delay, in a room with subdued light and with all windows tinted with a light-protective film. All sample treatment and analytical procedures were also carried out in this room.

#### Sample preparation and pre-treatment

Commonly consumed vegetables and fruits were purchased from markets and stalls. Samples were chosen from various groups of vegetables and fruits with differing characteristics. Edible portions of the foods were sizereduced in a blender and 2–10 g immediately weighed for analysis.

Sample pre-treatment procedures were essentially those of AOAC (Williams, 1984), except for the introduction of a saponification step. Preliminary studies carried out by the authors have shown that saponification was able to remove other pigments (mainly chlorophyll) from the food samples, which would otherwise interfere in the chromatography process, especially in the HPLC method. The saponification process did not appear to affect the  $\beta$ -carotene content, although there was some loss of lutein and other xanthophylls, which have no vitamin A activity.

To duplicate portions of the test sample were added a volume of 100% (w·v) potassium hydroxide equal to the weight of the food sample used, and 40 ml of ethanol. The mixture was saponified on an electric heating mantle for 30 min. The saponified mixture was cooled and extracted with 25 ml portions of hexane until the extract was colourless. The hexane extracts were pooled, washed until free of alkali, dried over sodium sulphate, and reduced to a small volume by heating over a water-bath with the aid of a stream of oxygen-free nitrogen. The resulting solution was made up immediately to a suitable volume (e.g. 25 ml) with hexane, referred to hereafter as the 'test solution'.

The total carotenoid concentration of the test solution was determined as described in the next section. The solution was next subjected to two chromatographic and subsequent quantitation procedures: (1) by open-column chromatography using magnesia and Hyflo Super Cel mixture and quantitation using absorbance reading at 450 nm (AOAC method); and (2) high-pressure liquid chromatography and detection and quantitation at



Fig. 2. UV Vis absorption spectra of (a) lycopene and  $\gamma$ -carotene in hexane (2 µg ml); (b) z- and  $\beta$ -carotene in hexane (2 µg/ml); (c) cryptoxanthin in hexane (2 µg ml); (d) zeaxanthin and lutein in ethanol (4 µg/ml).



Fig. 2—*contd.*(e)  $\beta$ -apo-carotenal in petroleum ether (2  $\mu$ g ml).

436 nm (HPLC method). The analytical procedures are summarised in Fig. 3.

# Determination of total carotenoid concentration

The test solution was read directly in a spectrophotometer at 450 nm and a  $\beta$ -carotene standard curve was used to calculate the total carotenoid content. The results obtained are referred to as having been obtained by the 'direct spectrophotometric method'.

The total carotenoid concentration was also determined by the highpressure liquid chromatographic (HPLC) method described below.

## **Open-column chromatography (AOAC method)**

A suitable volume (e.g. 10 ml) of the test solution was pipetted into a glass column prepacked with a mixture of activated magnesia (Sea Sorb 43) (Fisher Scientific Co. or Sigma Chemical Co.) and diatomaceous earth (Hyflo Super Cel) (Fisher Scientific Co.). in the ratio of 1:1, for chromatography using the AOAC method (Williams, 1984).  $\beta$ -Carotene was eluted from the column with approximately 80 ml of 10% (v/v) acetone in hexane. The eluate was evaporated on a water-bath with the aid of a stream of nitrogen and made up to a suitable volume (e.g. 10 ml) with hexane. The absorbance of the solution was read in a spectrophotometer at 450 nm and the concentration of  $\beta$ -carotene calculated by comparison against a calibration curve prepared with the  $\beta$ -carotene standard.



Fig. 3. Summary of analytical procedures.

# High-pressure liquid chromatography (HPLC method)

### HPLC conditions

A Waters high-pressure liquid chromatograph equipped with a 440 nm fixedwavelength detector was used. A 436 nm wavelength kit was fitted onto the detector and an attenuation of 0.02 absorbance units full scale (AUFS) was set. Another detector with a 313 nm wavelength kit and connected in series to the first, was found to be useful in assisting monitoring of carotenoids. particularly the *cis*-isomers of carotenes. A stainless steel 30 cm  $\times$  3.9 mm i.d. 10  $\mu$ m  $\mu$ Bondapak C<sub>18</sub> column was used for the chromatographic separation. This was preceded by a Waters Guard-PAK pre-column module housing a disposable Guard-PAK pre-column insert packed with the same material as that in the analytical column. Sample injection volumes, dispensed using a Rheodyne 7125 injector, were usually 50 to 100  $\mu$ l. A Waters 6000A solvent delivery system was used to deliver the mobile phase (acetonitrile-methanol–ethyl acetate, 88:10:2, v/v) at the rate of 2.0 ml/min. Peak areas were quantified with a Waters 730 Data Module.

# Chromatography of carotenoids

Hexane in the test solution was first evaporated off on a water-bath with the aid of nitrogen gas. The residue was immediately redissolved in a suitable volume of the mobile phase. After passing through a 0.45  $\mu$ m regenerated cellulose membrane filter, suitable volumes were injected into the chromatograph. Identification and quantitation of the carotenoids were carried out by comparing with reference carotenoids similarly chromatographed. Some food samples were found to contain a few carotenoids which could not be identified. The concentrations of these carotenoids were estimated as  $\beta$ -carotene. The concentrations of individual carotenoids were summed to give 'sum of carotenoids'. Total peak areas from the chromatograms were used to determine total carotenoid content, calculated using a  $\beta$ -carotene standard curve.

To assist in the identification of carotenoids in the samples, the pigments were eluted from the magnesia column (by the AOAC method) using a stepwise increase in the proportion of acetone in hexane. The absorption spectrum of each eluate was obtained and an aliquot injected into the HPLC to determine its purity and retention time. These data were compared with those given by authentic carotenoid standards. Details of these procedures were reported in Tee & Lim (1990).

### **RESULTS AND DISCUSSION**

This paper presents results for forty vegetables and fourteen fruits, listed in Tables 1 and 2, respectively. The English names of the foods are first listed, followed by the local names (in the Malay and other languages) and the scientific names. Where the English names are not available, the scientific names of the foods are used in the tables as well as the figures in this paper.

## Total carotenoid content

The total carotenoid concentration of each vegetable or fruit was determined by two methods: (1) taking the absorbance reading of the hexane extract at 450 nm previous to any chromatographic separation ('direct spectrophotometric method'); and (2) calculating the concentration from total peak area of HPLC chromatograms (HPLC method). There are two major limitations of total carotenoid concentrations determined by either method. Detection at a single wavelength (450 nm for the 'direct spectrophotometric method' and 436 nm for the HPLC method) is not adequate since absorption maxima for the various carotenoids differ considerably. The calculation based on  $\beta$ -carotene as the reference standard

 TABLE 1

 Names of Vegetables Studied

Green, leafy Cashew leavesPucuk gajusAnacardium occidentale Ceylon spinachRemayongBasella rubraC. griffithiiCemperaiChampereia griffithiiChinese cabbagePak-coyBrassica chimensisChinese cabbagePak-coyBrassica chimensisChinese chivesKucaiAllium odorumChinese kaleKai-lan-coyBrassica junceaCoriander leavesDaun ketumbarCoriandrum saticumCurry leavesKarrupillayMurray koenigiiDrumstick leavesDaun ketorMoringa oleiferaFern shootsPucuk pakuDiplazium esculentumH. javanicaPegaga gajahHydrocotyle javanicaLettuceSang-coyLactuca satiraMint leavesDaun melanMerinda actirifoliaN. citrifoliaDaun melanMerinda actirifoliaN. oleraceaTankiNeptunia oleraceaPapaya shootsDaun hetikCarica papayaSalted vegetableHum-coy—S. nigrumRantiSolaman nigrumSpinachBayam merahAmaranthus viridisSpinachBayam putihAmaranthus gangeticusSpring onionDaun havangAllium fistulosumSwamp cabbageKaacang buncisPhaseolus cularisSurgumKaacang buncisPhaseolus cularisSymanCekur manisSalted vegetableHum-coy——S. nigrumRantiSolama nigrumSolaman nigrumSpinachBayam putih<	English name	Local name	Scientific name
Cashew leavesPucuk gajusAnacardium occidentaleCyolon spinachRemayongBasella rubraC. griffthiiCemperaiChampereia griffthiiChinese cabbagePak-coyBrassica chimensisChinese chivesKucaiAllium odorumChinese chivesKucaiAllium odorumChinese kaleKai-lan-coyBrassica jimeeaCoriander icavesDaun ketumbarCoriandrum saticumCurry leavesKarrupillayMurray koenigiiDrumstick leavesDaun kelorMoringa olefferaFern shootsPucuk pakuDiplatium esculentumH. jaranicaPegaga gajahHydrocoryle jacanicaLettuceSang-coyLactuca saticaMint leavesDaun MengkuluMorinda citrifoliaN. oleraceaTankiNeptunia oleraceaPapaya shootsDaun hetikCarica papayaSaled vegetableHum-coy—S. androgynusCekur manisSauropus androgynusSesbaniaDaun turiSesbania grandiforaSpinachRaum putihAmaranthus gangeticusSpring onionDaun bavangAllium fistulosumSwamp cabbageKaueag baotoPaowangGreen, non-leafyCCascag baotoChill, greenLada hijauCapsicun annuamFour-angled beanKacang baotoPaowang Figna sinensisDottersCacang panjangFigna sinensisLog bean (dark green)Kacang baotoPaseolus culgarisLog bean (light green)	Green, leafy		
Ceylon spinachRemayongBasella rubraC. griffithiiComporaiChampereia griffithiiChinese cabbagePak-coyBrassica chinensisChinese cabbageKacaiAllium odorumChinese kaleKai-lan-coyBrassica alboglabraChinese kaleKai-lan-coyBrassica innecaCoriander ieavesDaan ketumbarCoriandrum satirumCurry leavesKarrupillayMurray KoenigiiDrumstick leavesDaan kelorMoringa oleiferaFern shootsPucuk pakuDiplazium esculeatumH. jaranicaPegaga gajahHydrocotyle jaranicaLettuceSang-coyLactuca satiraMint leavesDaan pudinaMentha arcensisM. cirrifoliaDaan betikCarica papayaSalted vegetableHum-coy—S. androgynusCekur manisSauropus androgynusSesbariiaDaan turiSesbania grandiforaS. nigrumRantiSolanum nigrumSpinach, redBayam merahAmaranthus ciridisSpinach, redBayam merahAmaranthus gangeticusSpring onionDaan bacangAlliun fistalosumSwamp cabbageKaag botorPsophocarpus tetragonolohusFrench beanKacang botorPsophocarpus tetragonolohusFrench beanKacang botorPsophocarpus tetragonolohusSynach, redBayam merahAmaranthus ciridisSpring onionDaun bacangAlliun fistalosumSwamp cabbageKaang botorPsophocarpus tetragono	Cashew leaves	Pucuk gajus	Anacardium occidentale
C. griffuhii Cemperal Champereia griffuhii Chinese cabbage Pak-coy Brassica chimensis Chinese chives Kucai Allium odorum Chinese Kale Kai-lan-coy Brassica ibnoglabra Chinese kale Kai-lan-coy Brassica ibnoglabra Chinese mustard leaves Sawi Brassica ibnoglabra Chinese mustard leaves Sawi Brassica ibnoglabra Curry leaves Daun ketumbar Coriandrum satirum Curry leaves Daun ketor Moringa oleifera Fern shoots Pucuk paku Diplazium esculentum H. jutanica Pegaga gajah Hydrocotyle javanica Lettuce Sang-coy Lactuca satira Mint leaves Daun pudina Mentha arcensis M. citrifolia Daun Mengkudu Morinda citrifolia N. oleracea Tanki Neptunia oleracea Papaya shoots Daun betik Carica papaya Salted vegetable Hum-coy — S. androgynus Cekur manis Sauropus androgynus Sesbania Daun turi Sesbania grandifora Spinach Bayam merah Amaranthus grangeticus Spring onion Daun bavang Allium fistulosum Symach Bayam merah Amaranthus gangeticus Spring onion Daun bavang Allium fistulosum Swamp cubbage Kangkung Ipomoca quatica Tapioca shoots Pucuk ubi kayu Mambat utilisima Wolfberry leaves Kau-kei-coy Lycium chinense Green, non-leafy Chilli, green Lada hijan Capsicum annuam Fuen-angled bean Kacang bator Psophocarpus tetragonolobus French bean Kacang panjang Vigna sinensis Long bean (light green) Kacang panjang Vigna s	Cevlon spinach	Remayong	Basella rubra
Chinese cabbagePak-coyBrassica chinensisChinese chivesKucaiAllium odorumChinese kaleKai-lan-coyBrassica alboglahraChinese mustard leavesSawiBrassica ijunceaCoriander icavesDaan ketumbarCoriandem sativumCurry leavesKarrupillayMurray koenigiiDrumstick leavesDaan kelorMoringa oleiferaFern shootsPucuk pakuDiplazium esculeatumH. jaranicaPegaga gajahHydrocotyle jaranicaLettuceSang-coyLactuca satiraMint leavesDaun MengkuduMorinda cirrifoliaN. oleraceaTankiNeptunia oleraceaPapaya shootsDaun turiSesbaniaSandrogynasCekur manisSauropus androgynusSesbaniaDaun turiSesbania grandiforaS. nigrumRantiSolanun nigrumSpinachBayam merahAmaranthus viridisSpinachBayam merahAmaranthus viridisSpinachBayam putihAmaranthus viridisSpinachBayam merahAmaranthus viridisSpinachBayam bayaManihot utilissimaWolfberry leavesKacang batorPsophocarpus tetragonolobusFrench beanKacang panjangVigna sinens	C. griffithii	Cemperai	Champereia griffithii
Chinese chivesKucaiAllium odorumChinese kaleKai-lan-coyBrassica alboglabraChinese mustard leavesSawiBrassica janceaCoriandern leavesDaun ketombarCoriandern saticumCurry leavesKarrupillayMurray koenigiiDrumstick leavesDaun kelorMoringa oleiferaFern shootsPucuk pakuDiplaziam esculentumH. jacanicaPegaga gajahHydrocotyle jacanicaLettuceSang-coyLactuca satiraMint leavesDaun pudinaMentha arcensisM. citrifoliaDaun betikCarica papyaN. oleraceaTankiNeptunia oleraceaPapaya shootsDaun betikCarica papyaSalted vegetableHum-coyS. androgynusCekur manisSauropus androgynusSesbaniaDaun turiSesbania grandiforaS. nigrumRantiSolanum ingrumSpinachBayam merahAmaranthus viridisSpinach, redBayam merahAmaranthus gageticusSpring onionDaun bawangAllium fistulosumSwamp cabbageKaacag buncisPhaseohus vulgarisLong bean (lark green)Kacang buncisPhaseohus vulgarisLong bean (light green)Kacang buncisPhaseohus vulgarisLong bean (light green)Kacang buncisPhaseohus vulgarisLong bean (light green)Kacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisLong bean (light green)Kacang	Chinese cabbage	Pak-cov	Brassica chinensis
Chinese kaleKai-lan-coyBrassica alboglabraChinese mustard leavesSawiBrassica junceaCoriander leavesDaan ketumbarCoriandrum saticumCurry leavesKarrupillayMurray koenigiiDrumstick leavesDaan kelorMoringa oleiferaFern shootsPucuk pakuDiplazium esculentumH. jaranicaPegaga gajahHydrocotyle jaranicaLettuceSang-coyLactuca sativaMint leavesDaun pudinaMentha arcensisM. citrifoliaDaun MengkuduMorinda citrifoliaN. oleraceaTankiNeptunia oleraceaPapaya shootsDaun betikCarica papayaSalted vegetableHum-coyS. androgynusCekur manisSauropus androgynusSesbaniaDaun turiSesbania grandiforaS. nigrumRantiSolanum nigrumSpinachBayam merahAmaranthus gangeticusSpring onionDaan bacangAllium fistulosumSwamp cabbageKangkungIpomoea aquaticaTapioca shootsPucuk ubi kayuManihot utilissimaWolfberry leavesKacang batorPsophocarpus tetragonolobusFrench beanKacang panjangVigna sinensisLong bean (dark green)Kacang panjangVigna sinensisLong bean (dark green)Ka	Chinese chives	Kucai	Allium odorum
Chinese mustard leavesSawiBrassica junceaCoriander leavesDaun ketnmbarCoriandrum satirumCurry leavesKarrupillayMurray koenigiiDrumstick leavesDaun ketorMoringa oleiferaFern shootsPucuk pakuDiplazium esculeatumH. jatanicaPegaga gajahHydrocotyle jatanicaLettuceSang-coyLactuca satiraMint leavesDaun mulinaMentha artensisM. citrifoliaDaun mengkuduMorinda citrifoliaN. oleraceaTankiNeptunia oleraceaPapaya shootsDaun betikCarica papayaSalted vegetableHum-coy—S. androgynusCekur manisSauropus androgynusSesbaniaDaun turiSesbania grandiforaS. nigrumRantiSolanum nigrumSpinachBayam merahAmaranthus gringeticusSpinachDaun bawangAllium fistulosumSwamp cabbageKangkungIpomoca aquaticaTapioca shootsPucuk ubi kayuManihot utilissimaWolfberry leavesKate-kei-coyLycium chinenseGreen, non-leafyCapsicum annuamChilli, greenLada hijauCapsicum annuamFour-angled beanKacang baniaPhaseolus tulgarisLong bean (dark green)Kacang panjangVigna sinensisLong bean (dark green)Kacang panjangVigna sinensisLong bean (dark green)Kacang panjangVigna sinensisPaprika Bell pepperLada hijau besarCapsicum annuam </td <td>Chinese kale</td> <td>Kai-lan-cov</td> <td>Brassica alboglahra</td>	Chinese kale	Kai-lan-cov	Brassica alboglahra
Coriander leavesDawn ketumbarCoriandrum saticumCurry leavesKarrupillayMurray koenigiiDrumstick leavesDawn kelorMoringa oleiferaFern shootsPucuk pakuDiplazium esculentumH. jaranicaPegaga gajahHydrocotyle jaranicaLettuceSang-coyLactuca satiraMint leavesDawn pudinaMentha arcensisM. cirrifoliaDawn MengkuduMorinda citrifoliaN. oleraceaTankiNeptunia oleraceaPapaya shootsDawn MengkuduMorinda citrifoliaSalted vegetableHum-coyS. androgynusCekur manisSauropus androgynusSesbaniaDawn turiSelsania growfiforaS. nigrumRantiSolanum nigrumSpinachBayam merahAmaranthus viridisSpinach, redBayam merahAmaranthus gangeticusSpring onionDawn bavangAllium fistulosumSwamp cabbageKangkungIpomoea aquaticaTapioca shootsPucuk ubi kayuManihot utilissimaWolfberry leavesKaacang botorPsophocarpus tetragonolobusFrench beanKacang botorPsophocarpus tetragonolobusFrench beanKacang panjangVigra sinensisLong bean (light green)Kacang panjangVigra sinensisLong bean (	Chinese mustard leaves	Sawi	Brassica juncea
Curry leavesKarrupillayMurray koenigiiDrumstick leavesDaun kelorMoringa oleiferaFern shootsPucuk pakuDiplazium esculeatumH. javanicaPegaga gajahHydrocotyle javanicaLettuceSang-coyLactuca sativaMint leavesDaun pudinaMentha arcensisM. citrifoliaDaun MengkuduMorinda citrifoliaN. oleraceaTankiNeptunia oleraceaPapaya shootsDaun tetikCarica papayaSalted vegetableHum-coyS. androgynusCckur manisSauropus androgynusSesbaniaDaun turiSesbania grandifioraS. nigrumRantiSolanum nigrumSpinachBayam putihAmaranthus viridisSpinach, redBayam merahAmaranthus gangeticusSpinach, redBayam merahAmaranthus gangeticusSpring onionDaun bawangAllium fistulosumSwamp cabbageKangkungIpomoca aquaticaTapioca shootsPucuk ubi kayuManihot utilissimaWolfberry leavesKacang botorPsophacarpus tetragonolobusFrench beanKacang botorPsophacarpus tetragonolobusFrench beanKacang panjangVigra sinensisLong bean (light green)Kacang panjangVigra sinensisPaprika Bell pepper <td>Coriander leaves</td> <td>Daun ketumbar</td> <td>Coriandrum sativum</td>	Coriander leaves	Daun ketumbar	Coriandrum sativum
Drumstick leavesDaun kelorMoringa oleiferaFern shootsPucuk pakuDiplazium esculentumH. jacanicaPegaga gajahHydrocotyle jacanicaLeitueeSang-coyLactuca sativaMint leavesDaun pudinaMentha arcensisM. citrifoliaDaun MengkuduMorinda citrifoliaN. oleraceaTankiNeptunia oleraceaPapaya shootsDaun betikCarica papayaSalted vegetableHum-coyS. androgynusCekur manisSauropus androgynusSesbaniaDaun turiSesbania grandifioraS. nigrumRantiSolanum nigrumSpinachBayam putihAmaranthus ciridisSpinach, redBayam merahAmaranthus gangeticusSpring onionDaun bawangAllium fistulosumSwamp cabbageKangkungIpomoea aquaticaTapioca shootsPucuk ubi kayuManihot utilissimaWolfberry leavesKacang buncisPhaseolus vulgarisLong bean (light green)Kacang buncisPhaseolus vulgarisLong bean (light green)Kacang panjangVigna sinensisPapita Bell peperLada hijau beaarCapsicum annuamSnake gourdKetola ularTricosanthes anguinaS. torcumTerung pipitSolanum torcumOthersCarrotLobak merahDaucus carotaChilli, redLada merahCapsicum annuamPumpkinLabu merahCapsicum annuamYam stalksBatang keladiHillium	Curry leaves	Karrupillay	Murray koenigii
Fern shootsPucuk pakuDiplazium esculentumH. javanicaPegaga gajahHydrocotyle javanicaLettuceSang-coyLactuca sativaMint leavesDaun pudinaMentha arcensisM. citrifoliaDaun MengkuduMorinda citrifoliaN. oleraceaTankiNeptunia oleraceaPapaya shootsDaun betikCarica papayaSalted vegetableHum-coyS. androgynusCekur manisSauropus androgynusSesbaniaDaun turiSesbania grandifioraS. nigrumRantiSolanum nigrumSpinachBayam putihAmaranthus viridisSpinach, redBayam putihAmaranthus gangeticusSpring onionDaun bawangAllium fistulosumSwamp cabbageKangkungIpomoea aquaticaTapioca shootsPucuk ubi kayuManihot utilissinaWolfberry leavesKacang botorPsophocarpus tetragonolobusFrench beanKacang batorPhaseolus vulgarisLong bean (light green)Kacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisPaprika Bell pepperLada hijau besarCapsicum annuan	Drumstick leaves	Daun kelor	Moringa oleifera
H. javanicaPegaga gajahHydrocotyle javanicaLettuceSang-coyLactuca sativaMint leavesDaun pudinaMentha arcensisM. citrifoliaDaun MengkuduMorinda citrifoliaN. oleraceaTankiNeptunia oleraceaPapaya shootsDaun betikCarica papayaSalted vegetableHum-coy—S. androgynusCekur manisSauropus androgynusSesbaniaDaun turiSeshania grandifioraSpinachBayam putihAmaranthus ciridisSpinachBayam merahAmaranthus ciridisSpinach, redBayam merahAmaranthus gangeticusSpring onionDaun kavangAllium fistulosumSwamp cabbageKaukei-coyLycium chinenseGreen, non-leafyChilli, greenLada hijauCapsicum annuamFrench beanKacang botorPsohocarpus tetragonolobusFrench beanKacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisPaprika Bell pepperLada hijau besarCapsicum annuamSnake gourdKetola ularTricosanthes anguinaS. torcunTerung pipitSolanum torcun <td>Fern shoots</td> <td>Pucuk paku</td> <td>Diplazium esculentum</td>	Fern shoots	Pucuk paku	Diplazium esculentum
LettuceSang-coyLactuca satiraMint leavesDaun pudinaMentha arcensisM. cirrifoliaDaun MengkuduMorinda citrifoliaN. oleraceaTankiNeptunia oleraceaPapaya shootsDaun betikCarica papayaSalted vegetableHum-coyS. androgynusCekur manisSauropus androgynusSesbaniaDaun turiSesbania grandifioraS. nigrumRantiSolanum nigrumSpinachBayam putihAmaranthus viridisSpinach, redBayam merahAmaranthus gangeticusSpring onionDaun bavcangAllium fistulosumSwamp cabbageKangkungIpomoca aquaticaTapioca shootsPucuk ubi kayuManihot utilissimaWolfberry leavesKacang botorPsophocurpus tetragonolobusFrench beanKacang botorPsophocurpus tetragonolobusFrench beanKacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisPaprika Bell pepperLada hijau besarCapsicum annuamSnake gourdKetola ularTricosanthes anguinaS. torvanTerang pipitSolanum torvamOthersCarrotLobak merahCaucus carotaChilli, redLada merahCapsicum annuamPumpkinLabu merahCucurbita maximaTomatoTomatoLycopersicum esculentiom	H. javanica	Pegaga gajah	Hydrocotyle javanica
Mint leavesDawn pudinaMentha arcensisM. citrifoliaDawn MengkuduMorinda citrifoliaN. oleraceaTankiNeptunia oleraceaPapaya shootsDawn betikCarica papayaSalted vegetableHum-coy—S. androgynusCekur manisSauropus androgynusSesbaniaDawn turiSesbania grandifioraS. nigrumRantiSolanum nigrumSpinachBayam putihAmaranthus viridisSpinach, redBayam merahAmaranthus gangeticusSpring onionDawn bawangAllium fistulosumSwamp cabbageKangkungIpomoea aquaticaTapioca shootsPucuk ubi kayuManihot utilissimaWolfberry leavesKacang botorPsophocarpus tetragonolobusFrench beanKacang botorPsophocarpus tetragonolobusFrench beanKacang panjangVigna sinensisLong bean (lark green)Kacang panjangVigna sinensisLong bean (lark green)Kacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisSolanumTricosanthes anguinaS. torrumStartaLada hijauCapsicum annuamSnake gourdKetola ularTricosanthes anguinaS. torrumTerung pipitSolanum torrumOthersCarrotLobak merahDaucus carotaChilli, redLada merahCapsicum annuamPumpkinLada merahCucurbita maximaTomatoTomatoLycopersicum esculentum <td>Lettuce</td> <td>Sang-cov</td> <td>Lactuca sativa</td>	Lettuce	Sang-cov	Lactuca sativa
M. citrifoliaDawn MengkuduMorinda citrifoliaN. oleraceaTankiNeptunia oleraceaPapaya shootsDaun betikCarica papayaSalted vegetableHum-coyS. androgynusCekur manisSauropus androgynusSesbaniaDaun turiSesbania grandifioraS. nigrumRantiSolanum nigrumSpinachBayam putihAmaranthus viridisSpinachBayam merahAmaranthus viridisSpinachBayam merahAmaranthus gangeticusSpring onionDaun bawangAllium fistulosumSwamp cabbageKangkungIpomoea aquaticaTapioca shootsPucuk ubi kayuManihot utilissimaWolfberry leavesKau-kei-coyLycium chinenseGreen, non-leafyCChilli, greenLada hijauCapsicum annuamFour-angled beanKacang botorPsophocarpus tetragonolobusFrench beanKacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisSolanum Treiung pipitSolanum torumStartaCarsturaTricosanthes anguinaS. torumTerung pipitSolanum torumOthersCarrotLobak merahCapsicum annuamFundaLada merahCapsicum annuamSnake gourdKetola ularTricosanthes anguinaS. torumTerung pipitSolanum torumOthersCarrotLobak merah	Mint leaves	Daun pudina	Mentha arvensis
N. oleraceaTankiNeptunia oleraceaPapaya shootsDaun betikCarica papayaSalted vegetableHum-coy—S. androgynusCekur manisSauropus androgynusSesbaniaDaun turiSesbania grandifforaS. nigrumRantiSolanum nigrumSpinachBayam putihAmaranthus viridisSpinach, redBayam merahAmaranthus gangeticusSpring onionDaun bawangAlliun fisulosuanSwamp cabbageKangkungIpomoea aquaticaTapioca shootsPucuk ubi kayuManihot utilissimaWolfberry leavesKacang butorPsophocarpus tetragonolobusFrench beanKacang butorPsophocarpus tetragonolobusFrench beanKacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisPapika Bell pepperLada hijauCapsicum annuumSnake gourdKetola ularTricosanthes anguinaS. torcumTerung pipitSolanum torcumOthersCarrotLobak merahCapsicum annuumPumpkinLada merahCapsicum annuumYam stalksBatang keladiYam stalks	M. citrifolia	Daun Mengkudu	Morinda citrifolia
Papaya shootsDaun betikCarica papayaSalted vegetableHum-coyS. androgynusCekur manisSauropus androgynusSesbaniaDaun turiSesbania grandifforaS. nigrumRantiSolanum nigrumSpinachBayam putihAmaranthus viridisSpinach, redBayam merahAmaranthus gangeticusSpring onionDaun bawangAllium fistulosumSwamp cabbageKangkungIpomoea aquaticaTapioca shootsPucuk ubi kayuManihot utilissimaWolfberry leavesKaarkei-coyLycium chinenseGreen, non-leafyChilli, greenLada hijauChilli, greenKacang batorPsophocarpus tetragonolobusFrench beanKacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisPapika Bell pepperLada hijauCapsicum annuumSnake gourdKetola ularTricosanthes anguinaS. torvanTerung pipitSolanum torvamOthersCarrotLobak merahCapsicum annuamPunpkinLabu merahCapsicum annuamPunpkinLabu merahCapsicum annuamYam stalksBatang keladiYam stalks	N. oleracea	Tanki	Neptunia oleracea
Salted vegetableHum-coy—S. androgynusCekur manisSauropus androgynusSesbaniaDaun turiSesbania grandifioraS. nigrumRantiSolanum nigrumSpinachBayam putihAmaranthus viridisSpinach, redBayam merahAmaranthus gangeticusSpring onionDaun bawangAllium fistulosumSwamp cabbageKangkungIpomoea aquaticaTapioca shootsPucuk ubi kayuManihot utilissimaWolfberry leavesKau-kei-coyLycium chinenseGreen, non-leafyChilli, greenLada hijauChilli, greenKacang botorPsophocarpus tetragonolobusFrench beanKacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisPaprika Bell pepperLada hijau besarCapsicum annuumSnake gourdKetola ularTricosanthes anguinaS. torvumTerung pipitSolanum torvumOthersCarrotLobak merahDaucus carotaChilli, redLada merahCapsicum annuumPumpkinLabu merahCucurbita maximaTomatoTomatoTomatoLycopersicum esculentum	Papaya shoots	Daun betik	Carica papaya
S. androgynusCekur manisSauropus androgynusSesbaniaDaun turiSesbania grandifioraS. nigrumRantiSolanum nigrumSpinachBavam putihAmaranthus viridisSpinach, redBayam merahAmaranthus gangeticusSpring onionDaun bawangAllium fistulosumSwamp cabbageKangkungIpomoea aquaticaTapioca shootsPucuk ubi kayuManihot utilissimaWolfberry leavesKau-kei-coyLycium chinenseGreen, non-leafyChilli, greenLada hijauChilli, greenLada hijauCapsicum annuumFour-angled beanKacang botorPsophocarpus tetragonolobusFrench beanKacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisDagrika Bell pepperLada hijau besarCapsicum annuumSnake gourdKetola ularTricosanthes anguinaS. torvumTerung pipitSolanum torvumOthersCarrotLobak merahDaucus carotaChilli, redLada merahCapsicum annuumPumpkinLabu merahCucurbita maximaTomatoTomatoTomatoLycopersicum esculentum	Salted vegetable	Hum-coy	
SesbaniaDawn turiSesbania grandifioraS. nigrumRantiSolanum nigrumSpinachBayam putihAmaranthus viridisSpinach, redBayam merahAmaranthus gangeticusSpring onionDawn bawangAllium fistulosumSwamp cabbageKangkungIpomoea aquaticaTapioca shootsPucuk ubi kayuManihot utilissimaWolfberry leavesKau-kei-coyLycium chinenseGreen, non-leafyCChilli, greenLada hijauCapsicum annuamFour-angled beanKacang botorPsophocarpus tetragonolobusFrench beanKacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisPaprika Bell pepperLada hijau besarCapsicum annuamSnake gourdKetola ularTricosanthes anguinaS. torvumTerung pipitSolanum torvumOthersCarrotLobak merahCapsicum annuamPumpkinLadu merahCucurbita maximaTomatoTomatoTomatoLycopersicum esculentum	S. androgynus	Cekur manis	Sauropus androgynus
S. nigrumRantiSolanum nigrumSpinachBayam putihAmaranthus viridisSpinach, redBayam merahAmaranthus gangeticusSpring onionDaun bawangAllium fistulosumSwamp cabbageKangkungIpomoea aquaticaTapioca shootsPucuk ubi kayuManihot utilissimaWolfberry leavesKau-kei-coyLycium chinenseGreen, non-leafyChilli, greenLada hijauChilli, greenKacang botorPsophocarpus tetragonolobusFrench beanKacang buncisPhaseolus vulgarisLong bean (lark green)Kacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisSnake gourdKetola ularTricosanthes anguinaS. torvumTerung pipitSolanum torvumOthersCarrotLobak merahCapsicum annuamPumpkinLabu merahCucurbita maximaTomatoTomatoLycopersicum esculentumYam stalksBotang keladiLycopersicum esculentum	Sesbania	Daun turi	Sesbania grandiflora
SpinachBayam putihAmaranthus viridisSpinach, redBayam merahAmaranthus gangeticusSpring onionDaan bawangAlliun fistulosumSwamp cabbageKangkungIpomoea aquaticaTapioca shootsPucuk ubi kayuManihot utilissimaWolfberry leavesKau-kei-coyLycium chinenseGreen, non-leafyCapsicum annuamFour-angled beanKacang botorPsophocarpus tetragonolobusFrench beanKacang buncisPhaseolus vulgarisLong bean (light green)Kacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisSnake gourdKetola ularTricosanthes anguinaS. torvumTerung pipitSolanum torvumOthersCarrotLobak merahChilli, redLada merahCapsicum annuamPumpkinLabu merahCucurbita maximaTomatoTomatoTomatoYam stalksBotang keladiLycopersicum esculentum	S. nigrum	Ranti	Solanum nigrum
Spinach, redBayam merahAmaranthus gangeticusSpring onionDaun bawangAllium fistulosumSwamp cabbageKangkungIpomoea aquaticaTapioca shootsPucuk ubi kayuManihot utilissimaWolfberry leavesKau-kei-coyLycium chinenseGreen, non-leafyCapsicum annuamFour-angled beanKacang botorPsophocarpus tetragonolobusFrench beanKacang buncisPhaseolus vulgarisLong bean (light green)Kacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisSonke gourdKetola ularTricosanthes anguinaS. torvumTerung pipitSolanum torvumOthersCarrotLobak merahCarrotLobak merahCapsicum annuamPumpkinLabu merahCucurbita maximaTomatoTomatoTomatoYam stalksBotang keladiLycopersicum esculentum	Spinach	Bayam putih	Amaranthus viridis
Spring onionDaan bawangAllium fistulosumSwamp cabbageKangkungIpomoea aquaticaTapioca shootsPucuk ubi kayuManihot utilissimaWolfberry leavesKau-kei-coyLycium chinenseGreen, non-leafyChilli, greenLada hijauCapsicum annuomFour-angled beanKacang botorPsophocarpus tetragonolobusFrench beanKacang buncisPhaseolus vulgarisLong bean (light green)Kacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisPaprika Bell pepperLada hijau besarCapsicum annuomSnake gourdKetola ularTricosanthes anguinaS. torvumTerung pipitSolanum torvumOthersCarrotLobak merahDaucus carotaChilli, redLada merahCapsicum annuomPumpkinLabu merahCucurbita maximaTomatoTomatoLycopersicum esculentumYam stalksBatang keladiLycopersicum esculentum	Spinach, red	Bayam merah	Amaranthus gangeticus
Swamp cabbageKangkungIpomoe aquaticaTapioca shootsPucuk ubi kayuManihot utilissimaWolfberry leavesKau-kei-coyLycium chinenseGreen, non-leafyChilli, greenLada hijauCapsicum annuamFour-angled beanKacang botorPsophocarpus tetragonolobusFrench beanKacang buncisPhaseolus vulgarisLong bean (dark green)Kacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisPaprika Bell pepperLada hijau besarCapsicum annuamSnake gourdKetola ularTricosanthes anguinaS. torvumTerung pipitSolanum torvumOthersCarrotLobak merahDaucus carotaChilli, redLada merahCapsicum annuamPumpkinLabu merahCucurbita maximaTomatoTomatoLycopersicum esculentumYam stalksBatang keladiLycopersicum esculentum	Spring onion	Daun bawang	Allium fistulosum
Tapioa shootsPucuk ubi kayuManihot utilissimaWolfberry leavesKau-kei-coyLycium chinenseGreen, non-leafyLada hijauCapsicum annuamFour-angled beanKacang botorPsophocarpus tetragonolobusFrench beanKacang buncisPhaseolus vulgarisLong bean (dark green)Kacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisPaprika Bell pepperLada hijau besarCapsicum annuamSnake gourdKetola ularTricosanthes anguinaS. torvumTerung pipitSolanum torvumOthersLobak merahCapsicum annuamPumpkinLabu merahCucurbita maximaTomatoTomatoTomatoLycopersicum esculentumYam stalksBatang keladiHang keladi	Swamp cabbage	Kangkung	Ipomoea aquatica
Wolfberry leavesKau-kei-coyLycium chinenseGreen, non-leafyChilli, greenLada hijauCapsicum annuamFour-angled beanKacang botorPsophocarpus tetragonolobusFrench beanKacang buncisPhaseolus vulgarisLong bean (dark green)Kacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisPaprika Bell pepperLada hijau besarCapsicum annuamSnake gourdKetola ularTricosanthes anguinaS. torvumTerung pipitSolanum torvumOthersLobak merahDaucus carotaChilli, redLada merahCapsicum annuamPumpkinLabu merahCucurbita maximaTomatoTomatoTomatoLycopersicum esculentumYam stalksBatang keladiFour states	Tapioca shoots	Pucuk ubi kayu	Manihot utilissima
Green, non-leafyLada hijauCapsicum annuamFour-angled beanKacang botorPsophocarpus tetragonolobusFrench beanKacang buncisPhaseolus vulgarisLong bean (dark green)Kacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisPaprika Bell pepperLada hijau besarCapsicum annuamSnake gourdKetola ularTricosanthes anguinaS. torvumTerung pipitSolanum torvumOthersCarrotLobak merahDaucus carotaChilli, redLada merahCapsicum annuamPumpkinLabu merahCucurbita maximaTomatoTomatoTomatoLycopersicum esculentumYam stalksBatang keladiLabu geladi	Wolfberry leaves	Kau-kei-coy	Lycium chinense
Chilli, greenLada hijauCapsicum annuamFour-angled beanKacang botorPsophocarpus tetragonolobusFrench beanKacang buncisPhaseolus vulgarisLong bean (dark green)Kacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisPaprika Bell pepperLada hijau besarCapsicum annuamSnake gourdKetola ularTricosanthes anguinaS. torvumTerung pipitSolanum torvumOthersCarrotLobak merahDaucus carotaChilli, redLada merahCapsicum annuamPumpkinLabu merahCucurbita maximaTomatoTomatoTomatoLycopersicum esculentumYam stalksBatang keladiSolanum torvum	Green, non-leafy		
Four-angled beanKacang botorPsophocarpus tetragonolobusFrench beanKacang buncisPhaseolus vulgarisLong bean (dark green)Kacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisPaprika Bell pepperLada hijau besarCapsicum annuamSnake gourdKetola ularTricosanthes anguinaS. torvumTerung pipitSolanum torvumOthersCarrotLobak merahDaucus carotaChilli, redLada merahCapsicum annuamPumpkinLabu merahCucurbita maximaTomatoTomatoTomatoLycopersicum esculentumYam stalksBatang keladiSolanu	Chilli, green	Lada hijau	Capsicum annuum
French beanKacang buncisPhaseolus vulgarisLong bean (dark green)Kacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisPaprika Bell pepperLada hijau besarCapsicum annuumSnake gourdKetola ularTricosanthes anguinaS. torvumTerung pipitSolanum torvumOthersLobak merahDaucus carotaChilli, redLada merahCapsicum annuumPumpkinLabu merahCucurbita maximaTomatoTomatoTomatoLycopersicum esculentumYam stalksBatang keladiSolanu	Four-angled bean	Kacang botor	Psophocarpus tetragonolobus
Long bean (dark green)Kacang panjangVigna sinensisLong bean (light green)Kacang panjangVigna sinensisPaprika Bell pepperLada hijau besarCapsicum annuumSnake gourdKetola ularTricosanthes anguinaS. torcumTerung pipitSolanum torcumOthersCarrotLobak merahDaucus carotaChilli, redLada merahCapsicum annuumPumpkinLabu merahCucurbita maximaTomatoTomatoLycopersicum esculentumYam stalksBatang keladiLabu	French bean	Kacang buncis	Phaseolus vulgaris
Long bean (light green)Kacang panjangVigna sinensisPaprika Bell pepperLada hijau besarCapsicum annuanSnake gourdKetola ularTricosanthes anguinaS. torcumTerung pipitSolanum torcumOthersCarrotLobak merahDaucus carotaChilli, redLada merahCapsicum annuamPumpkinLabu merahCucurbita maximaTomatoTomatoTomatoLycopersicum esculentumYam stalksBatang keladiSolanu	Long bean (dark green)	Kacang panjang	Vigna sinensis
Paprika Bell pepperLada hijau besarCapsicum annuumSnake gourdKetola ularTricosanthes anguinaS. torcumTerung pipitSolanum torcumOthersCarrotLobak merahDaucus carotaChilli, redLada merahCapsicum annuumPumpkinLabu merahCucurbita maximaTomatoTomatoLycopersicum esculentumYam stalksBatang keladi	Long bean (light green)	Kacang panjang	Vigna sinensis
Snake gourdKetola ularTricosanthes anguinaS. torvumTerung pipitSolanum torvumOthersCarrotLobak merahDaucus carotaChilli, redLada merahCapsicum annuumPumpkinLabu merahCucurbita maximaTomatoTomatoLycopersicum esculentumYam stalksBatang keladi	Paprika Bell pepper	Lada hijau besar	Capsicum annuum
S. torvumTerung pipitSolanum torvumOthersCarrotLobak merahDaucus carotaChilli, redLada merahCapsicum annuumPumpkinLabu merahCucurbita maximaTomatoTomatoLycopersicum esculentumYam stalksBatang keladi	Snake gourd	Ketola ular	Tricosanthes anguina
OthersLobak merahDaucus carotaCarrotLobak merahDaucus carotaChilli, redLada merahCapsicum annuunPumpkinLabu merahCucurbita maximaTomatoTomatoLycopersicum esculentumYam stalksBatang keladi	S. torvum	Terung pipit	Solanum torvum
CarrotLobak merahDaucus carotaChilli, redLada merahCapsicum annuumPumpkinLabu merahCucurbita maximaTomatoTomatoLycopersicum esculentumYam stalksBatang keladi	Others		
Chilli, redLada merahCapsicum annuunPumpkinLabu merahCucurbita maximaTomatoTomatoLycopersicum esculentumYam stalksBatang keladi	Carrot	Lobak merah	Daucus carota
PumpkinLabu merahCucurbita maximaTomatoTomatoLycopersicum esculentumYam stalksBatang keladi	Chilli, red	Lada merah	Capsicum annuum
TomatoTomatoLycopersicum esculentumYam stalksBatang keladi	Pumpkin	Labu merah	Cucurbita maxima
Yam stalks Botang keladi	Tomato	Tomato	Lycopersicum esculentum
	Yam stalks	Batang keladi	

	Names of Frans 5	ladea
English name	Local name	Scientific name
Banana (var. 1)	Pisang emas	Musa sapientum
Banana (var. 2)	Pisang tanduk	Musa sapientum
B. macrophylla	Buah kundang	Bouea macrophylla
Jackfruit	Nangka	Artocarpus heterophyllus
Mandarin orange	Limau Cina	Citrus reticulata
Mango (Black-gold)	Mangga	Mangifera indica
Musk lime	Limau kesturi	Citrus microcarpa
Orange	Limau manis	Citrus nobilis
Papaya	Betik	Carica papaya
Papaya exotica	Betik eksotika	Carica papaya
Plum, red	Buah plum	Prunus spp.
Starfruit Carambola	Belimbing manis	Averrhoa carambola
Tree tomato	Tomato pokok	Cyphomandra betacea
Watermelon, red	Tembikai	Citrullus vulgaris

 TABLE 2

 Names of Fruits Studied

is also an approximation since the different carotenoids have different extinction coefficients.

Tables 3 and 4 show total carotenoid concentrations in the vegetables and fruits obtained by the 'direct spectrophotometric' and HPLC methods. Figures 4 and 5 show the ratios of results obtained by the former method to those by the HPLC method, while Fig. 6 shows similar results obtained for the fruits. A ratio of unity indicates similar results were given by the two methods. For most of the vegetables and fruits (forty-six out of fifty-four samples studied), the ratios obtained were between 0.6 and 1.0, i.e. varying within  $\pm 20\%$  from 0.8. The results indicate that the HPLC method tended to give slightly higher total carotenoid concentrations. In spite of the limitations mentioned above, the data obtained could provide useful estimations of total carotenoid concentrations of vegetables and fruits.

#### **Carotenoid composition**

Only the HPLC method was able to give the carotenoid composition of the vegetables and fruits studied. The HPLC conditions employed gave satisfactory separation for lutein (retention time, RT = 3.6 min), cryptoxanthin (RT = 6.0 min), lycopene (RT = 7.5 min),  $\gamma$ -carotene (RT = 9.0 min),  $\alpha$ -carotene (RT = 10.2 min), and  $\beta$ -carotene (RT = 10.8 min). Zeaxanthin, structurally very similar to lutein (Fig. 1) was minimally separated from the latter.  $\beta$ -apo-8'-Carotenal, with the carbon skeleton shortened to 30, was also minimally separated from lutein. However, all three carotenoids had

# TABLE 3

Total Carotenoid Content<sup>a</sup> of Selected Vegetables Determined by Direct Spectrophotometric and HPLC Methods

Name of vegetable	Direct spectrophotometric method	HPLC method
Green, leafy		
Cashew leaves	1778	1 989
Cevlon spinach	3 4 3 2	4 5 5 6
C. griffithii	11833	17 310
Chinese cabbage	2 703	3728
Chinese chives	3 688	4 369
Chinese kale	4 395	5 299
Chinese mustard leaves	3 2 3 7	3741
Coriander leaves	4 0 5 0	4 3 4 4
Curry leaves	11 569	14170
Drumstick leaves	12986	14 565
Fern shoots	1944	2 278
H. javanica	4 281	4832
Lettuce	151	162
Mint leaves	5412	6 747
M citrifolia	7.047	1777
X oleracea	14 788	16 708
Papaya shoots	2120	2517
Salted vegetable	7 31 2	9635
S androgynus	31.877	39.910
Seshania	21.366	37.896
S nigrum	8 3 3 0	9.467
Spinach	5 481	7.058
Spinach red	6111	6 788
Spring onion	1885	1 5.17
Swamp cabbaga	2 () 2 2	2078
Tapiaga shoots	5 7 2 9	6075
Walthamin Lanax	5750	0975
wonderry leaves	10.376	12773
Green, non-leafy		
Chilli, green	678	810
Four-angled bean	555	589
French bean	525	784
Long bean (dark green)	776	1 070
Long bean (light green)	368	792
Paprika Bell pepper	300	453
Snake gourd	302	338
S. torvum	179	215
Others		
Carrot	9468	9 997
Chilli, red	5678	6 077
Pumpkin	1 830	2212
Tomato	1 276	1 224
Yam stalks	92	131

" Mean of duplicate analysis, expressed as  $\mu g$  per 100 g edible portion of sample.

TA	BL	Æ	4
			•

Total Carotenoid Content<sup>a</sup> of Selected Fruits Determined by Direct Spectrophotometric and HPLC Methods

Name of fruit	Direct spectrophotometric method	HPLC method
Banana (var. 1)	99	119
Banana (var. 2)	239	332
B. macrophylla	1 029	1 353
Jackfruit	193	197
Mandarin orange	669	916
Mango (Black-gold)	540	615
Musk lime	406	419
Orange	334	515
Papava	2 489	3 664
Papaya exotica	3 107	4 0 5 7
Plum, red	230	286
Starfruit Carambola	726	1 483
Tree tomato	1 604	1 583
Watermelon (red)	4 748	4 365

" Mean of duplicate analysis: expressed as  $\mu g$  per 100 g edible portion of sample.



Fig. 4. Ratio of total carotenoid concentrations of green, leafy vegetables determined by direct spectrophotometric and HPLC methods.



Fig. 5. Ratio of total carotenoid concentrations of green non-leafy and other vegetables determined by direct spectrophotometric and HPLC methods.

slightly different absorption spectra (Fig. 2). When vegetable extracts were fractionated on the magnesia column using step-wise increase of acetone in hexane as the eluant, the fraction eluted from the column with an RT of  $3\cdot3$  min in the HPLC chromatogram was found to have an absorption spectrum similar to that of lutein.  $\alpha$ - and  $\beta$ -Carotenes, differing only in the position of the double-bond in one of the two end groups (Fig. 1) were not completely separated. However, there was no difficulty in accurate identification and quantitation of these two pigments.



Fig. 6. Ratio of total carotenoid concentrations of fruits determined by direct spectrophotometric and HPLC methods.



**Fig. 7.** HPLC chromatogram of carotenoid standards. Detector 436 nm. 0·02 AUFS. Other chromatography conditions as given in text. Concentrations of lutein, cryptoxanthin and lycopene were 0·5  $\mu$ g ml, and of  $\alpha$ -,  $\beta$ - and  $\gamma$ -carotenes were 1·0  $\mu$ g ml. 100  $\mu$ l used for injection. 1 = Lutein: 2 = cryptoxanthin: 3 = lycopene; 4 =  $\gamma$ -carotene: 5 =  $\alpha$ -carotene: 6 =  $\beta$ -carotene.

A chromatogram of a mixture of these carotenoids is given in Fig. 7. It can be seen that elution order of the carotenoids on the reverse-phase  $C_{18}$ column was as expected, i.e. the more polar compounds were eluted earlier. As can be seen from the chromatogram, the oxygenated carotenoids or xanthophylls were eluted early. Lutein and zeaxanthin, the dihydroxy pigments, were eluted first, followed by the hydroxy carotenoid cryptoxanthin, and then the straight-chain carotenoid lycopene. The non-polar carotenoid hydrocarbons,  $\gamma$ -,  $\alpha$ - and  $\beta$ -carotenes were eluted last from the column.

The concentrations of the major carotenoids quantitated are given in Tables 5 and 6 for the vegetables and fruits, respectively. The carotenoids are tabulated in the order of their elution from the HPLC column, except for 'other carotenoids' not identified. Zero values in the tables refer to levels of carotenoids which were not detected or could not be quantitated accurately using the procedure as described. Figures 8–10 give the composition of the carotenoids, expressed as the percentage of each carotenoid to the sum of all carotenoids.

		Content <sup>a</sup> of	T. Major Caro	ABLE 5 tenoids in Selected	Vegetables			
Name of vegetable	Lutein	Cryptoxanthin	Lycopene	gamma-Carotene	alpha-Carotene	heta-Carotene	Others <sup>b</sup>	Sum <sup>c</sup>
Green, leafy								
Cashew leaves	773	0	0	0	0	1 342	0	2115
Ceylon spinach	1 299	0	0	c	0	3 533	0	4832
C. griffithii	9 871	0	0	0	3 677	3 218	0	16766
Chinese cabbage	963	0	0	0	0	3 0 2 2	0	3985
Chinese chives	1 083	0	0	0	0	3 509	0	4 592
Chinese kale	1 540	0	0	0	0	4092	0	5 631
Chinese mustard leaves	1019	0	0	0	0	2 928	0	3947
Coriander leaves	1 339	0	0	0	0	3 167	0	4 506
Curry leaves	5 252	0	0	0	0	9 328	0	14 579
Drumstick leaves	7128	0	0	0	0	7 536	0	14 663
Fern shoots	1 002	0	0	0	0	1438	0	2440
H. javanica	1 305	0	0	0	0	3840	0	5145
Lettuce	73	0	0	0	0	67	0	170
Mint leaves	1 701	0	0	0	0	4836	0	6 537
M. citrifolia	8 842	0	0	0	Û	3 108	1 296	13 246
N. oleracca	6 236	0	0	0	0	11 395	0	17631
Papaya shoots	821	0	0	0	0	1 829	0	2650
Salted vegetable	7318	0	0	0	0	3 395	С	10713
S. androgynus	29913	0	0	0	0	13351	3 292	46556
Sesbania	20.213	0	0	0	0	13611	5 890	39714
S. nigrum	2888	0	0	0	0	7 048	0	9936

# E-Siong Tee, Chin-Lam Lim

Spinach	4175	0	0	0	0	3177	0	7352
Spinach, red	2 ()47	0	0	0	0	5088	0	7134
Spring onion	323	0	0	0	0	1 282	0	1 605
Swamp cabbage	335	С	0	0	0	1 895	0	2 229
Tapioca shoots	1676	0	0	0	0	5720	0	7 396
Wolfberry leaves	7 591	0	0	0	0	5867	0	13458
Green. non-leafy								
Chilli, green	386	0	0	0	0	468	0	854
Four-angled bean	142	0	0	0	0	476	0	617
French bean	460	0	0	0	0	236	154	849
Long bean (dark green)	423	0	0	0	0	569	153	1 144
Long bean (light green)	300	0	0	0	0	412	146	857
Paprika Bell pepper	223	0	0	0	0	267	0	490
Snake gourd	225	0	0	0	0	148	0	372
S. torum	154	0	0	0	0	74	0	228
Others								
Carrot	0	0	0	0	3410	6 7 6 9	0	10179
Chilli, red	941	1 754	0	0	0	1 663	1971	6328
Pumpkin	94()	0	0	0	756	578	0	2273
Tomato	130	0	723	0	0	365	0	1218
Yam stalks	80	0	0	0	0	61	0	141

" Mean of duplicate analysis: expressed as  $\mu g$  per 100 g edible portion of sample.

<sup>b</sup> Unidentified carotenoids.
 <sup>c</sup> Summation of all carotenoids tabulated.

Name of fruit	Lutein	Cryptoxanthin	Ly coperate	gamma Carotene	alpha-Carotene	bena-Carotene	$Others^{h}$	Sum
Banana (var. 1)	27	0	. 0	0	62	40	c	128
Banana (var. 2)	37	0	0	()	157	<u>5</u> 6	0	286
B. macrophylla	457	155	0	52	0	301	<del>1</del> 15	1477
Jackfruit	95	17	0	0	С	56	56	223
Mandarin orange	113	688	С	59	0	81	151	1601
Mango (Black-gold)	С	0	0	C	0	615	0	615
Musk lime	65	446	С	0	0	<u>, 1</u>	0	522
Orange	30	332	0	0	0	35	218	605
Papaya	C	1483	2,003	118	0	228	767	4 125
Papaya exotica	0	615	2333	189	0	125	304	3 760
Plum, red	149	40	C	0	0	127	С	316
Starfruit Carambola	66	1 ()66	C	0	0	2.	551	1710
Tree tomato	0	1 236	0	0	0	500	C	1834
Watermelon (red)	0	157	5 301	<u>()</u> 6	0	324	0	6171

Content<sup>al</sup> of Major Carotenoids in Selected Fruits **TABLE 6** 

<sup>*a*</sup> Mean of duplicate analysis: expressed as  $\mu g$  per 100 g edible portion of sample. <sup>*b*</sup> Unidentified carotenoids.

<sup>e</sup> Summation of all carotenoids tabulated.



Fig. 8. Carotenoid composition of green, leafy vegetables.

For most of the green leafy vegetables, simple HPLC chromatograms were obtained and the major carotenoids detected were rather consistent (Table 5 and Fig. 8). In most cases, only  $\beta$ -carotene and lutein were obtained. The former was found in all the vegetables studied, and was clearly the major carotenoid in most of the vegetables. In twenty of the green leafy vegetables studied,  $\beta$ -carotene made up over 50% of the sum of all carotenoids



Fig. 9. Carotenoid composition of green, non-leafy and other vegetables.



Fig. 10. Carotenoid composition of fruits.

quantitated. For the remaining seven samples, at least 20% of the carotenoids was  $\beta$ -carotene. Lutein was also detected in all vegetables in fairly high proportions. Except for five samples, lutein made up over 25% of the sum of all carotenoids in these vegetables. The other carotenoids were encountered infrequently. x-Carotene was found only in *C. griffithii*, whilst  $\gamma$ -carotene. lycopene and cryptoxanthin were not detected under the conditions employed. For three of the vegetables in these two groups, a small proportion (<15%) of the carotenoids was contributed by a few unidentified carotenoids.

Carotenoid compositions of the green non-leafy vegetables were similar to those for the green leafy varieties (Table 5 and Fig. 9). Seven of the eight vegetables in the former group were found to have over 25% of lutein and over 30% of  $\beta$ -carotene. A few unidentified carotenoids were detected (constituting about 15% of the sum of carotenoids) in three vegetables; namely, French bean and the two varieties of long beans. These carotenoids appear to be characteristic of these leguminous vegetables.

In contrast to the green vegetables, the carotenoid composition of the other vegetables was rather different (Table 5 and Fig. 9). Although  $\beta$ -carotene and lutein were found in all these fruit and root vegetables, several other carotenoids were encountered.  $\alpha$ -Carotene was detected in carrot and pumpkin, while cryptoxanthin was found in red chilli. Lycopene was detected only in tomato, and made up about 60% of the carotenoids quantitated.

The fruits also presented rather different carotenoid compositions from

those obtained for the green vegetables (Table 6 and Fig. 10). There was no clear pattern of carotenoids present in the samples studied.  $\beta$ -Carotene was detected in all the fruits, but its proportion varied considerably, ranging from 100% in mango to less than 10% for seven other fruits. Lutein was found in nine of the fourteen fruits studied, but in smaller proportions than in green vegetables. Cryptoxanthin was found in most of the fruits studied, and contributed to over 50% of the carotenoids in five of the fruits. As for the vegetables,  $\alpha$ -carotene was infrequently encountered, having been detected only in the two banana species, contributing about 50% of total carotenoids. Lycopene also occurred infrequently, detected only in papaya (including the cultivar Exotica) and watermelon (red variety). In the last named, it constituted over 80% of all the carotenoids.  $\gamma$ -Carotene was found in small proportions (<5%) in five of the fruits studied. Seven of the fruits studied also had significant proportions of the unidentified carotenoids.

#### $\beta$ -Carotene content

 $\beta$ -Carotene concentrations determined by the AOAC and HPLC methods are tabulated in Tables 7 and 8. Figures 11–13 show the ratios of  $\beta$ -carotene determined by the two methods for the green leafy vegetables, green non-leafy and other vegetables and fruits, respectively.

For the green leafy vegetables, the ratios of  $\beta$ -carotene determined by the AOAC and HPLC methods clustered between 0.8–1.2, i.e. varying within



Fig. 11. Ratio of  $\beta$ -carotene and retinol equivalent (RE) of green, leafy vegetables determined by AOAC and HPLC methods.



Fig. 12. Ratio of  $\beta$ -carotene and retinol equivalent (RE) green, non-leafy and other vegetables determined by AOAC and HPLC methods.

 $\pm 20\%$  from unity (Table 7 and Fig. 11). Only in one vegetable was the ratio outside this range; *C. griffithii* was found to have an exceptionally high ratio of 2·1. This high ratio was due to the presence of  $\alpha$ -carotene in this vegetable which was eluted together with  $\beta$ -carotene from the magnesia column and erroneously estimated together with it in the AOAC method. This was confirmed by injecting the eluate from the magnesia column into the high-pressure liquid chromatograph and observing the two carotenes in the



Fig. 13. Ratio of  $\beta$ -carotene and retinol equivalent (RE) of fruits determined by AOAC and HPLC methods.

TAI	<b>BLE</b>	7
-----	------------	---

 $\beta$ -Carotene Content" of Selected Vegetables Determined by the AOAC and HPLC Methods

Name of vegetable	AOAC method	HPLC method
Green, leafy		
Cashew leaves	1 278	1 342
Cevlon spinach	2714	3 5 3 3
C. griffithii	6 788	3 218
Chinese cabbage	2 604	3 0 2 2
Chinese chives	2739	3 509
Chinese kale	3 500	4 0 9 2
Chinese mustard leaves	2 536	2928
Coriander leaves	2 745	3167
Curry leaves	8 241	9328
Drumstick leaves	7 724	7 536
Fern shoots	1 273	1 4 3 8
H. javanica	3 266	3 840
Lettuce	83	97
Mint leaves	4411	4836
M. citrifolia	3 0 7 4	3 108
N. oleracea	11 459	11 395
Papava shoots	1 709	1829
Salted vegetable	2 704	3 3 9 5
S. androgynus	12 363	13 351
Sesbania	12697	13611
S. nigrum	6 760	7 048
Spinach	2947	3177
Spinach, red	4 646	5 088
Spring onion	1 404	1 282
Swamp cabbage	1 729	1 895
Tapioca shoots	4 607	5720
Wolfberry leaves	6 4 1 4	5 867
Green, non-leafy		
Chilli, green	388	468
Four-angled bean	443	476
French bean	221	236
Long bean (dark green)	520	569
Long bean (light green)	285	412
Paprika Bell pepper	154	267
Snake gourd	138	148
S. toruun	82	74
Others		
Carrot	9 0 2 7	6 769
Chilli, red	1 743	1 663
Pumpkin	1 1 70	578
Tomato	352	365
Yam stalks	52	61

<sup>*a*</sup> Mean of duplicate analysis; expressed as  $\mu$ g per 100 g edible portion of sample.

Name of fruit	AOAC method	HPLC method
Banana (var. 1)	82	40
Banana (var. 2)	219	92
B. macrophylla	303	301
Jackfruit	55	56
Mandarin orange	120	81
Mango (Black-gold)	495	615
Musk lime	18	12
Orange	40	25
Papaya	208	228
Papava exotica	314	321
Plum, red	113	127
Starfruit Carambola	77	28
Tree tomato	582	599
Watermelon (red)	246	324

**TABLE 8** β-Carotene Content" of Selected Fruits Determined by the AOAC and HPLC Methods

" Mean of duplicate analysis; expressed as  $\mu g$  per 100 g edible portion of sample.

chromatogram. For most of the other vegetables, the ratio tended to be less than 1.0, indicating that the HPLC method gave slightly higher results.

The ratios of  $\beta$ -carotene content determined by the AOAC and HPLC methods for the green non-leafy and other vegetables were more varied (Table 7 and Fig. 12). However, most of the ratios were within the narrow range of 0.8 to 1.2. For pumpkin, the ratio was exceptionally high at 2.1, due to the presence of  $\alpha$ -carotene which was giving erroneously high results by the AOAC method.

For the fruits, there was considerable variation in the ratios of  $\beta$ -carotene concentration given by the two methods (Table 8 and Fig. 13). For half of the fruits studied, the ratios were between 0.8 and 1.2. For the only two fruits with  $\alpha$ -carotene, i.e. the two species of banana, the ratios were greater than 2.0. The reason for this over-estimation by the AOAC method has been explained above. Three other fruits with ratios of about 1.5 were musk lime, orange and starfruit. These fruits were found to have low levels of  $\beta$ -carotene (less than 100  $\mu$ g per 100 g edible portion), which made up only a small proportion of all the carotenoids detected. The relatively insensitive and non-specific nature of the AOAC method, especially for foods with low  $\beta$ -carotene, could be the reason for the over-estimation by this method.

Results obtained from the HPLC method showed that highest concentrations of  $\beta$ -carotene were found in green leafy vegetables, particularly *S. androg vnus*, sesbania and *N. oleracea* (Table 7). All these three

local vegetables had a  $\beta$ -carotene content of over 11000  $\mu$ g per 100 g of edible portion. Six other green leafy vegetables were found to have a  $\beta$ -carotene content from 5000-10000  $\mu$ g per 100 g of vegetable. With the exception of carrot ( $\beta$ -carotene about 9000  $\mu$ g per 100 g), none of the green non-leafy and other vegetables was found to have high  $\beta$ -carotene content. All the fruits studied were not rich in  $\beta$ -carotene, with concentrations of less than 1000  $\mu$ g per 100 g of sample. The sample of mango studied was interesting in that all the carotenoids were solely  $\beta$ -carotene, and the concentration was the highest of all the fruits.

### Vitamin A activity (retinol equivalent)

Conventionally, the nutritional significance of carotenoids is related to the pro-vitamin A activity of these compounds. Although over 500 carotenoids have been reported to occur naturally, only a few are known to possess provitamin A activity and occur in significant amounts in natural foods (Bauernfeind, 1972; Goodwin, 1986). For vitamin A activity, a carotenoid must have at least one unsubstituted  $\beta$ -ionone ring with an attached polyene side chain of at least eleven carbon atoms. Consistent with these important structural requirements, the following carotenoids identified in this study have been known to possess pro-vitamin A activity:  $\beta$ -carotene,  $\alpha$ -carotene,  $\gamma$ -carotene, and cryptoxanthin. The vitamin A activity of  $\beta$ -carotene, expressed as  $\mu g$  retinol equivalent (RE) was calculated as RE = ( $\mu g \beta$ -carotene) 6 (NAS, 1980). The other three carotenoids mentioned, possessing only one unsubstituted  $\beta$ -ionone ring, may be expected to have about 50% of the biological activity of  $\beta$ -carotene. The formula used for these provitamin A carotenoids was therefore RE = ( $\mu g$  carotenoid)/12.

The RE values calculated from  $\beta$ -carotene determined by the HPLC method and those calculated based on all carotenoids with pro-vitamin A activity by the same analytical method are given in Table 9 for the vegetables. The differences in RE between the two methods of calculation, expressed as a percentage of RE from all carotenoids, are also given in the table. For the green leafy vegetables, with the exception of *C. griffithii*, since no other carotenoids with pro-vitamin A activity were detected, there was no difference in RE calculated by the two methods. For *C. griffithii*, excluding  $\alpha$ -carotene from the calculation would result in a 36% error in RE content for the vegetables. For the other vegetables; namely, carrot, red chilli and pumpkin, differences in RE ranged from 20 to 40%.

Since other pro-vitamin A carotenoids besides  $\beta$ -carotene were detected in all the fruits studied except mango, the RE values calculated on the basis of all these carotenoids were higher than those calculated based on  $\beta$ -

#### TABLE 9

Retinol Equivalent (RE)<sup>a</sup> of Selected Vegetables Determined by the AOAC and HPLC Methods

Name of vegetable	AOAC method <sup>b</sup>	· · · · · · · · · · · · · · · · · · ·	HPLC me	thod
		$\beta$ -carotene <sup>c</sup>	total <sup>d</sup>	<sup>o</sup> a difference <sup>e</sup>
Green, leafy	· • • • • • • • • • • • • • • • • • • •			
Cashew leaves	213	224	224	0
Ceylon spinach	452	589	589	0
C. griffithii	1 1 3 1	536	843	36.4
Chinese cabbage	434	504	504	()
Chinese chives	457	585	585	0
Chinese kale	583	682	682	0
Chinese mustard leaves	423	488	488	0
Coriander leaves	457	528	528	0
Curry leaves	1 374	1 555	1 5 5 5	Õ
Drumstick leaves	1.287	1 2 5 6	1 256	0
Fern shoots	212	240	240	0
H. javanica	544	640	640	0
Lettuce	14	16	16	0
Mint leaves	735	806	806	0
M. citrifolia	512	518	518	0
N. oferacea	1910	1 899	1 899	0
Papaya shoots	285	305	305	õ
Salted vegetable	451	566	566	õ
S. androgynus	2 061	2 225	2 2 2 5	0
Sesbania	2116	2 269	2 269	0
S. nigrum	1127	1175	1175	0
Spinach	491	530	530	Õ
Spinach, red	774	848	848	0
Spring onion	234	214	214	0
Swamp cabbage	288	316	316	0
Tapioca shoots	768	953	953	0
Wolfberry leaves	1 069	978	978	0
Green, non-leafy				
Chilli. green	65	78	78	0
Four-angled bean	74	79	79	0
French bean	37	39	39	0
Long bean (dark green)	87	95	95	0
Long bean (light green)	47	69	69	0
Paprika Bell pepper	26	45	45	0
Snake gourd	23	25	25	0
S. torvum	14	12	12	0
Others				
Carrot	1 504	1 1 2 8	1412	20-1
Chilli, red	290	277	423	34.5
Pumpkin	195	96	159	39.5
Tomato	59	61	61	0
Yam stalks	9	10	10	0

<sup>*a*</sup> Mean of duplicate analysis; expressed as  $\mu g$  per 100 g edible portion of sample. <sup>*b*</sup> Calculated as RE = ( $\mu g$  carotene) 6. <sup>*b*</sup> Based on  $\beta$ -carotene only; RE = ( $\mu g \beta$ -carotene) 6. <sup>*d*</sup> Based on  $\beta$ -carotene and all other pro-vitamin A carotenoids, i.e.  $\alpha$ -carotene,  $\gamma$ --carotene. cryptoxanthin:  $RE = [(\mu g \beta \text{-carotene}) 6] + [(\mu g \text{ other carotenoids}) 12].$ 

<sup>e</sup> Calculated as:	$\begin{bmatrix} \text{total } RE - RE \text{ from } \beta \text{-carotene} \\ \text{total } RE \end{bmatrix}$	× 100.
-----------------------------	--	--------

TABLE	E 10
-------	------

Retinol Equivalent (RE)<sup>4</sup> of Selected Fruits Determined by the AOAC and HPLC Methods

Name of fruit	AO 4C method <sup>b</sup>	HPLC method		
		$\beta$ -carotene <sup>c</sup>	total <sup>4</sup>	vo difference <sup>e</sup>
Banana (var. 1)	1.4	7	12	43.8
Banana (yar. 2)	36	15	28	46.0
B. macrophylla	51	50	67	25.6
Jackfruit	9	9	11	1.2-8
Mandarin orange	20	14	76	82.2
Mango (Black-gold)	82	103	103	0
Musk lime	3	2	39	95-1
Orange	7	4	32	87.1
Papaya	35	38	171	77-8
Papaya exotica	52	54	120	55.6
Plum, red	19	21	24	13.7
Starfruit Carambola	13	5	93	95.1
Tree tomato	97	100	203	50.8
Watermelon (red)	41	54	99	45.8

" Mean of duplicate analysis; expressed as  $\mu g$  per 100 g edible portion of sample.

<sup>b</sup> Calculated as  $RE = (\mu g \text{ carotene})$  6.

<sup>4</sup> Based on  $\beta$ -carotene only: RE = ( $\mu g \beta$ -carotene) 6.

<sup>*d*</sup> Based on  $\beta$ -carotene and all other pro-vitamin A carotenoids, i.e.  $\alpha$ -carotene,  $\gamma$ -carotene, cryptoxanthin: RE = [( $\mu g \beta$ -carotene) 6] + [( $\mu g \beta$  other carotenoids) 12].

<sup>c</sup> Calculated as  $\begin{bmatrix} \text{total } \mathbf{RE} - \mathbf{RE} \text{ from } \beta \text{-carotene} \\ \hline \text{total } \mathbf{RE} \end{bmatrix} \times 100.$ 

carotene only (Table 10). The vitamin A activity calculated on the basis of  $\beta$ -carotene alone would result in RE values which were 12–87% lower.

The RE values calculated from carotene determined by the AOAC method and all carotenoids with pro-vitamin A activity by the HPLC method are given in Table 9 for the vegetables. As can be expected, with the exception of *C. griffithii*, the ratios of RE calculated by the two methods were exactly the same as ratios obtained for  $\beta$ -carotene (Fig. 11), because  $\beta$ -carotene was the only pro-vitamin A carotenoid detected for these vegetables. Owing to the presence of  $\alpha$ -carotene in *C. griffithii*, the RE for this vegetable was falsely elevated when calculated from carotene determined by the AOAC method since  $\alpha$ -carotene possesses only half the pro-vitamin A activity of  $\beta$ -carotene.

Similarly, for the green non-leafy vegetables, since no other pro-vitamin A carotenoids besides  $\beta$ -carotene were detected, the ratios of RE calculated from carotene determined by the AOAC method and all carotenoids with pro-vitamin A activity by the HPLC methods were exactly the same as the



Fig. 14. Classification of vegetables according to retinol equivalent (RE).

ratios obtained for  $\beta$ -carotene (Table 9 and Fig. 12). With the exception of two vegetables, the ratios were found to be between 0.8 and 1.2.

For three of the other vegetables studied, the ratios for RE were between this range (Fig. 8). The ratio for pumpkin was exceptionally high at about 2·0, indicating a falsely elevated RE given by the AOAC method. The reason for this is as given for *C. griffithii* above. On the other hand, RE for red chilli given by the AOAC method would be falsely low since the procedure was not able to quantitate cryptoxanthin.

Similarly, because of the presence of other pro-vitamin A carotenoids in fruits (except for mango), the ratios for RE determined by the AOAC method to that by the HPLC method were different from the  $\beta$ -carotene ratios. Ratios for RE were all lower than those for  $\beta$ -carotene, and were generally below 1.0 (Table 10 and Fig. 13). These data show that the AOAC method was underestimating the vitamin A activity of the fruits. since the method was not able to quantitate the other pro-vitamin A carotenoids.



Fig. 15. Classification of fruits according to retinol equivalent (RE).

To facilitate easy identification of vegetables which are good sources of vitamin A activity, these foods were grouped into four categories; namely, low ( $<100 \mu g$  RE per 100 g edible portion), medium ( $100-499 \mu g$  RE), high ( $500-999 \mu g$  RE) and very high ( $>1000 \mu g$  RE) (Fig. 14). Vegetables with high and very high RE were all green leafy vegetables, with the exception of carrot, a root vegetable. Of particular interest are four local vegetables with over 1000  $\mu g$  RE; namely, sesbania, *S. androgynus*, *N. oleracea*, and *S. nigrum*. Several other local vegetables were in the high RE category. A few other green leafy vegetables and red chilli and pumpkin made up the group with medium RE. All the green non-leafy vegetables, as well as yam stalk and tomato were found to be poor sources of vitamin A.

A similar grouping was made for the fruits studied (Fig. 15). None of the fruits may be considered as having high or very high vitamin A activity. The two species of papaya, tree tomato and mango were found to have medium RE, while the other fruits were poor sources of vitamin A, particularly the first variety of banana and jackfruit.

#### CONCLUSION

A non-aqueous reverse-phase HPLC method for the determination of carotenoids in various vegetables and fruits has been developed. The method uses the basic configuration of an HPLC system, and would thus be useful for routine determinations. A ternary mixture of acetonitrile, methanol and ethyl acetate was used to separate the carotenoids isocratically in an octadecylsilane ( $C_{18}$ ) column. A fixed wavelength detector at 436 nm was used to detect the carotenoids, the peaks being monitored and quantitated in an integrator. The method gave satisfactory separation and quantitation of lutein, cryptoxanthin, lycopene,  $\gamma$ -,  $\alpha$ - and  $\beta$ -carotenes. The emphasis has

been on major carotenoids that occur in sufficient amounts to contribute significantly to dietary intake.

This is the first report on the concentration of major carotenoids in relation to total carotenoids in a number of Malaysian vegetables and fruits. It is also the first report of a parallel study of carotenoid determination by the AOAC and the HPLC methods for a fairly large number of samples. Findings from the study have clearly shown that the HPLC method would give a more complete picture of the carotenoid composition as well as a more accurate quantitation of the pro-vitamin A activity of vegetables and fruits. Depending on the composition of the carotenoids present, the AOAC method could under- or over-estimate the  $\beta$ -carotene concentration and. therefore, the RE activity. The HPLC procedure reported could be useful for updating the vitamin A activity of plant materials in the current Malaysian Food Composition Table, thereby providing the correct identification of foods rich in pro-vitamin A activity. The nutritional significance of the findings is clear since these foods are important sources of vitamin A for the majority of the communities in the country. Furthermore, there is currently a great deal of interest in carotenoids not possessing vitamin A activity as they may be associated with lower cancer risk.

#### ACKNOWLEDGEMENTS

The authors thank Dr M. Jegathesan, Director of the Institute for Medical Research, for his encouragement and support of the project and granting permission to publish this paper. Thanks are also conveyed to Dr Y. H. Chong of Palm Oil Research Institute of Malaysia for his valuable comments on the manuscript. The technical assistance of Ms Rajamanohari and Ms S. C. Khor is also acknowledged. The authors also thank Messrs F. Hoffmann La-Roche of Switzerland for providing several reference carotenoids which were not obtainable commercially. The project was funded under the Intensification of Research Priority Areas (IRPA) Programme of the Ministry of Science, Technology and Environment Malaysia (Project No. IMR 53).

## REFERENCES

- Bauernfeind, J. C. (1972). Carotenoid vitamin A precursors and analogs in foods and feeds. J. Agric. Food Chem., 20, 456–73.
- Beecher, G. R. & Khachik, F. (1984). Evaluation of vitamin A and carotenoid data in food composition tables. J. Nat. Cancer Inst., 73, 1397–404.

- Bureau, J. L. & Bushway, R. J. (1986). HPLC determination of carotenoids in fruits and vegetables in the United States. J. Food Sci., 51, 128–30.
- De Ritter, E. (1981). Absorption maxima and E (1%/1 cm) values for carotenoids. In *Carotenoids as Colorants and Vitamin A Precursors*. ed. J. C. Bauernfeind, Academic Press, New York, pp. 883–923.
- Goodwin, T. W. (1986). Metabolism, nutrition and function of carotenoids. *Annu. Rev. Nutr.*, **6**, 273–97.
- NAS (1980). Fat-soluble vitamins: Vitamin A. In *Recommended Dictary Allowances*, (9th edn). National Academy of Sciences, Washington, DC, pp. 55-60.
- Olson, J. A. (1986). Carotenoids, vitamin A and cancer. J. Nutr., 116, 1127-30.
- Peto, R., Doll, R., Buckley, J. D. & Sporn, M. B. (1981). Can dietary beta-carotene materially reduce human cancer rates? *Nature*, **290**, 201–8.
- Tee, F. S. (1988). *Carotenoids and Retinoids in Human Nutrition*, ed. C. L. Lim & Y. H. Chong, Institute for Medical Research, Kuala Lumpur.
- Tee, E. S. & Lim, C. L. (1990). Development of a HPLC method for the analysis of carotenoids in vegetables and fruits. Paper presented at the seminar on Advances in Food Research III, Serdang, Malaysia, 19–20 November, 1990.
- Tee, E. S. & Lim, C. L. (1991). The Analysis of Carotenoids and Retinoids: A Review. Food Chemistry, **41**(2), 147–93.
- Tee, E. S., Mohd. Ismail, N., Mohd. Nasir, A. & Khatijah, I. (1988). Nutrient Composition of Malaysian Foods. ASEAN Sub-Committee on Protein: Food Habits Research and Development. Kuala Lumpur.
- Temple, N. J. & Basu, T. K. (1988). Does beta-carotene prevent cancer? A critical appraisal. Nutr. Res., 8, 685–701.
- Underwood, B. A. (1984). Vitamin A in animal and human nutrition. In *The Retinoids*. Volume 1. ed. M. B. Sporn, A. B. Roberts & D. S. Goodman, Academic Press, Orlando, pp. 281–392.
- Williams, S. (ed). (1984). *Official Methods of Analysis of the AOAC*, (14th edn), Association of Official Analytical Chemists, Virginia, pp. 834–5.
- Zakaria, M., Simpson, K., Brown, P. R. & Krstulovic, A. (1979). Use of reversedphase high-performance liquid chromatographic analysis for the determination of provitamin A carotenes in tomatoes. J. Chromatogr., 176, 109–17.